

DEFORMATION OF THE SADDLE POINT FROM HEAVY ION INDUCED FISSION FRAGMENT ANGULAR DISTRIBUTIONS

K. Mahata and S. Kailas

Nuclear Physics Division, Bhabha Atomic Research Centre, Mumbai 400 085, India

Determining the potential energy landscape in the deformation space continues to be a challenging problem [1]. There has been a long standing discrepancy [2] between the deduced effective moment of inertia (\mathfrak{I}_{eff}), which is related to the saddle point shape, from the analysis of α induced fission reactions and the values predicted by the models. Though the agreement between the rotating finite range model (RFRM) [3] predictions and the experimental results (see Fig. 1) is good for high ($\frac{Z^2}{A}$), the deduced \mathfrak{I}_{eff} values are more close to rotating liquid drop model (RLDM) [4] predictions for low ($\frac{Z^2}{A}$). There were no experimental data for intermediate values of ($\frac{Z^2}{A}$). Further, these models are not well tested in heavy-ion induced fusion-fission reactions involving large angular momentum values.

In the present work, the \mathfrak{I}_{eff} values are obtained from detailed statistical model analysis of the measured fission fragment angular distributions for $^{19}\text{F} + ^{188,192}\text{Os}, ^{194,198}\text{Pt}$ systems. The multichance nature of fission has been taken into account explicitly and the calculation is constrained by fitting the measured evaporation residues and fission cross-sections along with the prefission neutron multiplicity data from systematics. The existing literature data have been also reanalyzed to cover a wide range of ($\frac{Z^2}{A}$). Though the values of the extracted \mathfrak{I}_{eff} are consistently larger than those predicted by the RFRM, their trend is found to agree with that of the RFRM over the whole range of ($\frac{Z^2}{A}$) studied (see Fig. 1, X-axis is correct for finite spin of the fissioning nuclei). These results will provide a useful constraint for different models, which are used to predict the fission barriers and the shapes for shell stabilized super heavy nuclei.

- [1] Peter Möller, Arnold J. Sierk and Akira Iwamoto, Phys. Rev. Lett **92**, 072501 (2004).
- [2] R. Vandenbosch, *Proc. Beijing International Symposium on physics at tandem*, p355 (1986), World Scientific.
- [3] A. J. Sierk, Phys. Rev. **C 33**, 2039 (1986).
- [4] S. Cohen, F. Plasil and W.J. Swiatecki, Ann. Phys. (N.Y.) **82**, 557 (1974).

